



PERSONS WITH A HISTORY OF ALCOHOL OR OTHER DRUG ABUSE

- Problem drinkers were shown in one study to be 23 times more likely to have a gambling problem than persons without a drinking problem (Welte et al., 2001).
- Pathological or problem gamblers had 7 times the rate of alcohol dependence than nongamblers and low-risk gamblers (NORC, 1999).
- A 1999 study (Pasternak & Fleming, 1999) revealed that almost one in every three persons with a gambling problem also abused alcohol (compared to about one in ten without gambling problems), and about twice as many problem gamblers used tobacco.
- In a study of over 21,000 high-school students, (Proimos, DuRant, Pierce, & Goodman, 1998) regular cocaine use and anabolic steroids were significantly associated with a report of problems connected to gambling.

PERSONS WITH A HISTORY OF MENTAL HEALTH ISSUES

- Individuals with concurrent psychiatric problems display much higher rates of disordered gambling than either adolescents or adults sampled from the general population (Shaffer et al., 1997).
- There are strong associations between pathological gambling and depression (Becona, Del Carmen Lorenzo, & Fuentes, 1996).
- High rates of personality disorders (e.g., obsessive-compulsive, avoidant, schizotypal and paranoid) are noted in the research (NORC, 1999).

ETHNIC MINORITIES

- Minority populations most likely have much higher rates of pathological gambling than Caucasians (Moore, Jadlos, & Carlson, 2000, as cited in Moore, 2001).
- Minorities spend about two and a half times more on gambling in a typical month than Caucasians (Moore, 2001a).
- Ethnic minority groups (except Asians) are over-represented among individuals classified as problem gamblers (Moore, 2001).

OLDER ADULTS

- The rate of problem gambling among Oregonian adults ages 65 and over is 1.2%, lower than any other age group in Oregon (Moore, 2001b); however, older adults who play at gambling facilities are six times more likely to be problem gamblers than other older adults randomly chosen from the same community, and 3.7 times as likely to be probable pathological gamblers (McNeilly & Burke, 2000).
- Older adults are considered at increased risk due to factors unique to that population, such as loneliness, isolation, physical or mental illness; additionally, in one study of older adults, the most likely reported motivations to gamble were: relaxation, boredom, passing time, and getting away for the day (McNeilly & Burke, 2000).
- Compared to those older adults surveyed from senior centers, older adults surveyed from bingo events had higher South Oaks Gambling Screen (SOGS) scores and greater gambling frequency and expenditures (Ladd, Molina, Kerins, & Petry, 2003).
- Among adults over 65 years of age, gambling is the most frequently identified social activity (McNeilly & Burke, 2001).
- From 1975 to 1997, the number of American adults 65 years of age or older that had ever gambled increased by 45% (NORC, 1999).
- From 1994 to 1998, the percentage of women 65 years of age or older who had ever gambled rose from by 20% (NORC, 1999).
- Some older adults may have cognitive impairment that interferes with their ability to make sound decisions.
- Many older adults may not understand addiction, making them less likely to identify a gambling problem; additionally, older adults appear less willing to seek assistance for a gambling problem than younger adults.
- There is also concern that gambling-related problems may be more difficult to identify because of increased isolation from co-workers, friends and family.